Over the age of 65 with A-Fib and not on an Anticoagulant? You may be eligible to join a Pioneering Clinical Trial!



Testing Abelacimab, an Investigational, Monthly Injectable for Stroke Prevention in A-Fib Patients



Goals of the Study

The LILAC - TIMI 76 trial is looking for participants who have non-valvular atrial fibrillation (AF), including paroxysmal AF, and have been told by their doctors that they can't take oral blood-thinning medication because they have a high risk of bleeding.

Who May Participate?



History of atrial fibrillation Current episode not necessary

Not on a current blood thinner like Eliquis®, Xarelto®, or Warfarin in the past 60 days

Be able to explain why you are unable to be on an oral blood thinner





If left atrial appendage closure (Watchman Device) isn't right for you, consider the LILAC - TIMI 76 trial

AFIB and Stroke

preventable cause of stroke is AF



X stroke risk



0/ patients require

life-long anticoagulation

If interested, please contact the HCL Research Team via: Phone: 020 7535 8311 Email: hcl.trials@nhs.net

*67% of patients with newly diagnosed AF in 2020 were not started on an anticoagulant within 12 months of diagnosis (Ko et al. (2022) JAMA). Reasons you may not be prescribed an anticoagulant include: Older age, fall risk, dementia, frailty, anemia, prior hip/pelvic fracture, chronic kidney disease, or personal choice (LILAC Poster V2 12Oct2023)